

CIALIS

If you are having any problems with your erection, then try Cialis, which is an effective impotency drug. With a single pill of Cialis, you can have immense sexual pleasure with your partner.

Drug usage

Cialis works by relaxing the muscles in the penis region. Once the muscles are relaxed, more blood flows into the region, which means better erection. It also helps in the erection to last for longer hours.



How taken

Never buy Cialis on your own. Always consult the doctor before buying Cialis. Care should be taken that you do not take larger amounts of Cialis. The drug can be taken with or without food. The drug can be taken 30 minutes ahead of any sexual activity. Only one Cialis pill has to be taken a day and you should have to note that there should be a gap of 24 hours between two dosages.



Missed dose/overdose

No one can miss a dose of Cialis as those who love to have that extra pleasure with their partner only use it. In case you have taken more of the drug, you may come across various symptoms like nausea, chest pain, irregular heartbeat, light-headedness and fainting. If you come across any of these overdose symptoms, never delay in calling the doctor.



More Information

Once your doctor has decided to provide you Cialis, you should never hide from him any of your allergies and also your health problems. Discuss with your physician if you have any of the following health issues: heart disease, recent heart attack, heart rhythm problems, stroke, congestive heart failure, angina, high/low pressure, Kidney/liver issues, blood cell problems, bleeding disorders, retinitis pigmentosa and Peyronie's disease.



Warnings

Never take Cialis if you are using any nitrate filled medicines as it can lead to a decrease of the blood pressure to dangerous levels. While taking Cialis, there are chances that you may have problems with your vision. Some people have experienced sudden vision loss but it is not yet known if Cialis is the real reason for the sudden vision loss. Those people who experienced such a condition were those who were diabetic or had high blood pressure or had vision problems earlier.



While taking Cialis, avoid alcohol as it can only increase the risk of side effects. You should also give up grape fruits and grape juice while taking Cialis.

Side effects

The allergic reactions that come with Cialis include hives, swelling (lips, tongue, mouth, throat) and breathing difficulty.



While using Cialis, you may also come across serious side effects including sudden vision and hearing loss, ringing in ears, chest pain, pain spreading to arms or shoulders, sweating, ill feeling, nausea, irregular heartbeat and seizures. If you find any of these side effects, stop Cialis and consult the doctor.

Some people may experience longer erections that may last for more than four hours. In this case, you have to contact the doctor immediately as longer erections can damage the penis.

The lesser-known side effects include stuffy nose, warmth in face/neck/chest, sore throat, diarrhoea, headache, upset stomach, back and muscles pains.

www.cialiscanada.org